

TABLE 4 - TABEL 4
 PROVISIONS RELATING TO SEED AND SEED SAMPLES - BEPALINGS BETREFFENDE SAAD EN SAADMONSTERS
 (*No. of footnote/*No. van voetnota)

Kind of plant		Maximum content (%)			Minimum percentage		Minimum number per weight (kg)	Prepacked seed		Exempted from indication "Prepacked seed"		Maximum mass (kg) of a seed lot
Botanical name	Common name	Other matter	Other seed	Weed seed	Germination by number	Viability	Germination by weighed replicate	Max.mass (g) per container	Approx. no. of seed per container	Max. mass (g) per container	Approx. no. of seed per container	
1	2	3	4	5	6	7	8	9	10	11	12	13
<i>×Agrotriticum</i> Cif. & Giacom	Agrotriticum	4	0,2		60			5 000	165 000	500	16 000	20 000
<i>Allium cepa</i> L.	Onion	4	0,2		60			500	170 500	50	17 050	10 000
<i>Allium fistulosum</i> L.	Bunching onion	4	0,2		60			500		50		10 000
<i>Allium fistulosum</i> L. × <i>A. cepa</i> L.	New Bunching onion	4	0,2		60			500		50		10 000
<i>Allium porrum</i> L.	Leek	4	0,2		60			500	198 000	50	19 800	10 000
<i>Anthephora pubescens</i> Nees (*7)	Bottle brush grass, (i) Uncoated seed (ii) Coated seed	10	0,3		20			5 000	905 000	500	90 500	10 000
		5	0,3		20			5 000	145 000	500	14 500	10 000
<i>Arachis hypogaea</i> L.	Groundnut	2	0,1		70			5 000	5 000 15 000	500	500 1 500	30 000
<i>Asparagus officinalis</i> L.	Asparagus	4	0,2		60			500	25 300	50	2 500	20 000
<i>Avena nuda</i> L. (*6)	Naked oats	0,5	0,3	TR	80			5 000	200 000	500	20 000	30 000
<i>Avena sativa</i> L. [incl. <i>A. byzantina</i> K. Koch]	Oats, Red oats	1	0,3	0,1	80			5 000	200 000	500	20 000	30 000
<i>Avena strigosa</i> Schreb.	Black oats	2,5	2,0	0,1	70			5 000		500		30 000
<i>Beta vulgaris</i> L.	Fodder beet, Garden beet, Sugar beet and Swiss chard	6	0,3		60			1 000	58 000	100	5 800	20 000
<i>Brassica napus</i> L. var. <i>napus</i>	Oilseed rape	2,0	0,2		70			500		50		10 000
<i>Brassica napus</i> L. var. <i>napobrassica</i> (L.) Rchb.	Forage rape and Swede	4	0,2		60			500	157 500	50	15 750	10 000
<i>Brassica oleracea</i> L.	Broccoli, Brussels sprouts, Cabbage, Cauliflower, Fodder kale, Kale, Kohlrabi and Savoy cabbage	4	0,2		60			500	157 500	50	15 750	10 000

Kind of plant		Maximum content (%)			Minimum percentage		Minimum number per weight (kg)	Prepacked seed		Exempted from indication "Prepacked seed"		Maximum mass (kg) of a seed lot
Botanical name	Common name	Other matter	Other seed	Weed seed	Germination by number	Viability	Germination by weighed replicate	Max.mass (g) per container	Approx. no. of seed per container	Max. mass (g) per container	Approx. no. of seed per container	
1	2	3	4	5	6	7	8	9	10	11	12	13
<i>Brassica rapa</i> L. [incl. <i>B. campestris</i> L.; <i>B. chinensis</i> L.; <i>B. pekinensis</i> (Lour.) Rupr.; <i>B. perviridis</i> (L.H. Bailey) L.H. Bailey]	Chinese cabbage, Pak- choi, Turnip	4	0,2		60			500	316 500	50	31 650	10 000
<i>Bromus catharticus</i> Vahl	Rescue grass	8	0,5		50			5 000	720 000	500	72 000	10 000
<i>Cajanus cajan</i> (L.) Huth	Pigeon pea	4	0,2		60							20 000
<i>Capsicum</i> spp.	Chilli, Hot pepper, Sweet pepper, Paprika and Pepper rootstock	4	0,2		60			500	83 500	50	8 350	10 000
<i>Cenchrus ciliaris</i> L.	Blue buffalo grass (i) Uncoated seed (ii) Coated seed	4	0,3		20			5 000	2 675 000	500	267 000	10 000
		4	0,3		20			5 000	290 000	500	29 000	10 000
<i>Chloris gayana</i> Kunth	Rhodes grass (i) Uncoated seed (ii) Coated seed	15	1,5	1,0	20	20	800 000	5 000	23 620 000	500	2 362 000	10 000
		5	1,5	1,0	20	20		5 000	2 580 000	500	258 000	10 000
<i>Citrullus lanatus</i> (Thunb.) Matsum. et Nakai	Watermelon	4	0,1		60			1 000	11 000	100	1 100	20 000
<i>Coriandrum sativum</i> L.	Coriander	1	0,3		60							10 000
<i>Cucumis melo</i> L.	Sweet melon	4	0,1		60			1 000	45 000	100	4 500	10 000
<i>Cucumis sativus</i> L.	Cucumber	4	0,1		60			1 000	38 000	100	3 800	10 000
<i>Cucumis</i> spp.	Cucumis rootstock	4	0,1		60			1 000		100		10 000
<i>Cucurbita maxima</i> Duch.	Pumpkin, Squash	4	0,1		60			1 000	5 000	100	500	20 000
<i>Cucurbita moschata</i> Duch.	Pumpkin, Squash	4	0,1		60			1 000	14 000	100	1 400	10 000
<i>Cucurbita pepo</i> L.	Squash, Vegetable marrow, Zucchini	4	0,1		60			1 000	14 000	100	1 400	20 000
<i>Cucurbita</i> spp.	Rootstock	4	0,1		60			1 000		100		10 000
<i>Dactylis glomerata</i> L.	Cocksfoot	15	2,5	0,5	60			5 000	8 280 000	500	828 000	10 000
<i>Daucus carota</i> L.	Carrot	4	0,2		60			500	413 000	50	41 300	10 000
<i>Desmodium intortum</i> (Mill.) Urb. (*2)	Green leaf desmodium	6	1,0	0,5	50			5 000	3 240 000	500	324 000	10 000
<i>Desmodium uncinatum</i> (Jacq.) DC. (*2)	Silver leaf desmodium	6	1,0	0,5	50			5 000	1 010 000	500	101 000	20 000
<i>Digitaria eriantha</i> Steud. [syn. <i>D. smutsii</i> Stent; <i>D.</i> <i>decumbens</i> Stent]	Smuts finger grass (i) Uncoated seed (ii) Coated seed	8,0	1,0	0,5	15			5 000	10 660 000	500	1 066 000	10 000
		8,0	1,0	0,5	15			5 000	5 235 000	500	523 500	10 000
<i>Echinochloa crus-galli</i> (L.) P. Beauv.	Japanese millet	10	0,7		60							10 000
<i>Eleusine corocana</i> (L.) Gaertn.	Finger millet	10	??		60							10 000
<i>Eragrostis curvula</i> (Schrad.) Nees (*5)	Weeping lovegrass	4	1,0	0,3	70			500	1 641 000	50	164 100	10 000

Kind of plant		Maximum content (%)			Minimum percentage		Minimum number per weight (kg)	Prepacked seed		Exempted from indication "Prepacked seed"		Maximum mass (kg) of a seed lot
Botanical name	Common name	Other matter	Other seed	Weed seed	Germination by number	Viability	Germination by weighed replicate	Max. mass (g) per container	Approx. no. of seed per container	Max. mass (g) per container	Approx. no. of seed per container	
1	2	3	4	5	6	7	8	9	10	11	12	13
<i>Eragrostis tef</i> (Zucc.) Trotter	Teff	4	1,0	0,3	70			500	2 304 000	50	230 400	10 000
<i>Fagopyrum esculentum</i> Moench	Buckwheat	6	??		60							10 000
<i>Festuca arundinacea</i> Schreb.	Tall fescue	4	1,5	0,3	60			5 000	2 500 000	500	250 000	10 000
× <i>Festulolium</i> Asch. & Graebn.	Festulolium	4	1,5	0,3	60			5 000		500		10 000
<i>Foeniculum vulgare</i> Mill.	Fennel	2	0,5		60							10 000
<i>Glycine max</i> (L.) Merrill	Soya bean	4	0,1		70			10 000	60 000 130 000	1 000	6 000 13 000	30 000
<i>Gossypium hirsutum</i> L.	Cotton	4	0,2		70			10 000	80 000	1 000	8 000	25 000
<i>Helianthus annuus</i> L.	Sunflower	4	0,1		70			10 000	160 000	1 000	16 000	25 000
<i>Hordeum vulgare</i> L. subsp. <i>vulgare</i> (*6)	Barley	2,5	0,3	TR	80			5 000	150 000	500	15 000	30 000
<i>Kummerowia striata</i> (Thunb.) Schindl. [syn. <i>Lespedeza striata</i> (Thunb.) Hook. & Arn.] (*1)	Common Lespedeza	4	0,3		60			5 000	3 750 000	500	375 000	10 000
<i>Lablab purpureus</i> (L.) Sweet [syn. <i>Dolichos lablab</i> L.]	Lab-Lab bean	6	??		60							20 000
<i>Lactuca sativa</i> L.	Lettuce	6	0,2		60			500	500 000	50	50 000	10 000
<i>Lespedeza cuneata</i> (Dum. Cours.) G. Don (*1)	Chinese Lespedeza	4	0,3		60			5 000	4 100 000	500	410 000	10 000
<i>Lespedeza striata</i> – see <i>Kummerowia striata</i>												
<i>Lolium multiflorum</i> Lam.	Italian and Westerwold ryegrass	4	1,5	0,3	60	60		5 000	2 500 000	500	250 000	10 000
<i>Lolium perenne</i> L.	Perennial ryegrass	4	1,5	0,3	60	60		5 000	2 500 000	500	250 000	10 000
<i>Lolium rigidum</i> Gaudin	Wimmeria ryegrass, Annual ryegrass	4	1,5	0,3	60	60		5 000	1 035 000	500	103 500	10 000
<i>Lolium xboucheanum</i> – see <i>Lolium xhybridum</i>												
<i>Lolium xhybridum</i> Hausskn. [syn. <i>L. xboucheanum</i> Kunth]	Hybrid ryegrass	4	1,5	0,3	60			5 000	1 925 000	500	192 500	10 000
<i>Lotus corniculatus</i> L. (*3)	Birdsfoot trefoil	4	0,6	0,3	60			5 000	4 070 000	500	407 000	10 000
<i>Lupinus albus</i> L. (*3)	White lupin	2	0,3	0,1	70			10 000	70 000	1 000	7 000	30 000
<i>Lupinus angustifolius</i> L. (*3)	Narrow leaf lupin	2	0,3	0,1	70			10 000	70 000	1 000	7 000	30 000
<i>Lupinus luteus</i> L. (*3)	Yellow lupin	2	0,3	0,1	70			10 000	90 000	1 000	9 000	30 000
<i>Lycopersicon esculentum</i> (syn. <i>L. lycopersicum</i>) – see <i>Solanum lycopersicum</i>												

Kind of plant		Maximum content (%)			Minimum percentage		Minimum number per weight (kg)	Prepacked seed		Exempted from indication "Prepacked seed"		Maximum mass (kg) of a seed lot
Botanical name	Common name	Other matter	Other seed	Weed seed	Germination by number	Viability	Germination by weighed replicate	Max. mass (g) per container	Approx. no. of seed per container	Max. mass (g) per container	Approx. no. of seed per container	
1	2	3	4	5	6	7	8	9	10	11	12	13
<i>Medicago littoralis</i> Rhode ex Loisel. (*2)	Strand medic	4	0,2		60			5 000	1 805 000	500	180 500	10 000
<i>Medicago polymorpha</i> L. (*2)	Burr medic	3	0,8	0,3	60			5 000	1 380 000	500	138 000	10 000
<i>Medicago rugosa</i> Desr. (*2)	Gama medic	3	0,5	0,3	60			5 000	705 000	500	70 500	10 000
<i>Medicago sativa</i> L. [incl. <i>Medicago x varia</i> Martyn; <i>M. falcata</i> L.] (*2)	Lucerne	3	0,5	0,3	70			2 000	730 000	200	73 000	10 000
<i>Medicago scutellata</i> (L.) Mill. (*2)	Snail medic	3	0,5	0,3	60			5 000	255 000	500	25 500	10 000
<i>Medicago truncatula</i> Gaertn. (*2)	Barrel medic	3	1,0	0,3	60			5 000	1 190 000	500	119 000	10 000
<i>Megathyrsus maximus</i> (Jacq.) B.K. Simon & S.W.L. Jacobs [syn. <i>Panicum maximum</i> Jacq.]	White buffalo grass, (i) Uncoated seed	2,5	0,7	0,5	10	20		5 000	11 035 000	500	1 103 500	10 000
	(ii) Coated seed	2,5	0,7	0,5	10	20		5 000	3 845 000	500	384 500	10 000
<i>Melilotus albus</i> Medik.	White sweet clover	6			60							10 000
<i>Nicotiana tabacum</i> L.	Tobacco	4	0,2		80			5 000	78 125 000	500	7 812 500	10 000
<i>Ornithopus compressus</i> L. (*1)	Yellow serradella	4	1,5	0,5	70			5 000	1 035 000	500	103 500	10 000
<i>Ornithopus sativus</i> Brot. (*3)	Serradella	2	0,5	0,3	60			5 000	1 135 000	500	113 500	10 000
<i>Oryza sativa</i> L.	Rice	4	0,1		70			10 000	660 000	1 000	66 000	30 000
<i>Panicum coloratum</i> L.	Small Buffalo grass (i) Uncoated seed	2	0,7	0,5	20			5 000		500		10 000
	(ii) Coated seed	2	0,7	0,5	20			5 000		500		10 000
<i>Panicum maximum</i> – see <i>Megathyrsus maximus</i>												
<i>Paspalum dilatatum</i> Poir.	Dallis grass	40	0,3		40			5 000	2 960 000	50	29 600	10 000
<i>Paspalum notatum</i> Flügge	Bahia	40	0,3		40			5 000	1 830 000	50	18 300	10 000
<i>Pastinaca sativa</i> L.	Parsnip	7	0,2		50			500	214 500	50	21 450	10 000
<i>Pennisetum clandestinum</i> Hochst. ex Chiov.	Kikuyu	5	0,3		60			500	185 500	50	18 550	10 000
<i>Pennisetum glaucum</i> (L.) R. Br.	Pearl millet	7	0,2		60			5 000	970 000	500	97 000	10 000
<i>Petroselinum crispum</i> (Mill.) Fuss	Parsley	6	0,2		50			500	324 000	50	32 400	10 000
<i>Phalaris aquatica</i> L.	Harding grass	6	0,3		60			5 000	2 980 000	500	298 000	10 000
<i>Phaseolus coccineus</i> L.	Kidney bean	4	0,1		60			5 000	5 000	500	500	30 000
<i>Phaseolus vulgaris</i> L.	Dry bean	4	0,1		60			5 000	20 000	500	2 000	30 000
<i>Phaseolus vulgaris</i> L.	Garden bean – Runner	4	0,1		60			5 000	20 000	500	2 000	30 000
<i>Phaseolus vulgaris</i> L.	Garden bean - Dwarf	4	0,1		60			5 000	20 000	500	2 000	30 000
<i>Pisum sativum</i> L. <i>sensu lato</i>	Dry pea, Garden pea	4	0,1		60			5 000	25 000	500	2 500	30 000
<i>Raphanus sativus</i> L.	Fodder radish	4	0,2		60			5 000	375 000	500	37 500	10 000
<i>Raphanus sativus</i> L.	Garden radish	4	0,2		60			500	60 000	50	6 000	10 000
<i>Ricinus communis</i> L.	Castor oil	4	0,1		70			10 000	50 000	1 000	5 000	20 000
<i>Secale cereale</i> L.	Rye	2	0,5	0,2	70			5 000	200 000	500	20 000	30 000
<i>Setaria sphacelata</i> (Schumach.) Stapf & C.E. Hubb.	Setaria	40	0,3		30			5 000	7 825 000	500	782 500	10 000

Kind of plant		Maximum content (%)			Minimum percentage		Minimum number per weight (kg)	Prepacked seed		Exempted from indication "Prepacked seed"		Maximum mass (kg) of a seed lot
Botanical name	Common name	Other matter	Other seed	Weed seed	Germination by number	Viability	Germination by weighed replicate	Max.mass (g) per container	Approx. no. of seed per container	Max. mass (g) per container	Approx. no. of seed per container	
1	2	3	4	5	6	7	8	9	10	11	12	13
<i>Sinapis alba</i> L.	White mustard	4	0,2		60			500	284 500	50	28 450	10 000
<i>Solanum lycopersicum</i> L. [syn. <i>Lycopersicon esculentum</i> Mill., <i>L. lycopersicum</i> (L.) H. Karst.]	Tomato	2	0,2		60			500	202 500	50	20 250	10 000
<i>Solanum melongena</i> L.	Aubergine, Brinjal, Eggplant	4	0,2		60			500	114 000	50	11 400	10 000
<i>Solanum</i> spp. (excluding <i>S. elaeagnifolium</i> Cav.)	Rootstock	4	0,2		60			500		50		10 000
<i>Sorghum bicolor</i> (L.) Moench subsp. <i>bicolor</i> [syn. <i>S. bicolor</i> (L.) Moench; <i>S. vulgare</i> Pers.; including <i>S. dochna</i> (Forssk.) Snowden]	Grain sorghum	4	0,1		70			5 000	150 000	500	15 000	30 000
<i>Sorghum</i> spp. (excl. <i>S. halepense</i> (L.) Pers.) (<i>S. bicolor</i> (L.) Moench subsp. <i>bicolor</i> × <i>S. bicolor</i> (L.) Moench subsp. <i>drummondii</i> (Steud.) de Wet ex Davidse [syn. <i>Sorghum sudanense</i> (Piper) Stapf] and hybrids with seeds not retained in tight bracts)	Forage sorghum	7	0,2		60			5 000	650 000	500	65 000	30 000
<i>Sorghum</i> spp. (excl. <i>S. halepense</i> (L.) Pers.) (<i>S. bicolor</i> (L.) Moench subsp. <i>drummondii</i> (Steud.) de Wet ex Davidse [syn. <i>S. sudanense</i> (Piper) Stapf] and hybrids with seeds mostly retained in tight bracts) (*4)	Forage sorghum	7	0,2		60	70		5 000	650 000	500	65 000	10 000
<i>Sorghum</i> spp. (excl. <i>S. halepense</i> (L.) Pers.) (<i>S. xalmum</i> Parodi) (*4)	Forage sorghum	7	0,2		60	70		5 000	650 000	500	65 000	30 000
<i>Sporobolus fimbriatus</i> (Trin.) Nees (*7)	Dropseed grass	4	1	0,3	70			500		50		10 000
<i>Stylosanthes hamata</i> (L.) Taub. (*1)	Caribbean stylo	10	1,0	0,5	50			5 000	2 450 000	500	245 000	10 000
<i>Trifolium fragiferum</i> L. (*3)	Strawberry clover	6	0,3		60			1 000	635 000	100	63 500	10 000
<i>Trifolium hirtum</i> All. (*3)	Rose clover	4	0,3		60			1 000	358 000	100	35 800	10 000
<i>Trifolium incarnatum</i> L. (*3)	Crimson clover	4	0,3		60			1 000	330 000	100	33 000	10 000
<i>Trifolium pratense</i> L. (*3)	Red clover	6	0,3		60			1 000	600 000	100	60 000	10 000
<i>Trifolium repens</i> L. (*2)	White clover	6	0,3		60			1 000	1 500 000	100	150 000	10 000

Kind of plant		Maximum content (%)			Minimum percentage		Minimum number per weight (kg)	Prepacked seed		Exempted from indication "Prepacked seed"		Maximum mass (kg) of a seed lot
Botanical name	Common name	Other matter	Other seed	Weed seed	Germination by number	Viability	Germination by weighed replicate	Max. mass (g) per container	Approx. no. of seed per container	Max. mass (g) per container	Approx. no. of seed per container	
1	2	3	4	5	6	7	8	9	10	11	12	13
<i>Trifolium resupinatum</i> L. (*3)	Persian clover	4	0,3		60			1 000	1 416 000	100	141 600	10 000
<i>Trifolium subterraneum</i> L. (*3)	Subterranean clover	6	0,3		60			1 000	119 000	100	11 900	10 000
<i>Trifolium vesiculosum</i> Savi (*1)	Arrow leaf clover	4	0,3		60			1 000	699 000	100	59 900	10 000
x <i>Triticosecale</i> Wittm. ex A. Camus	Triticale (<i>Triticum</i> x <i>Secale</i>)	3	0,5		80			5 000	130 000	500	13 000	30 000
<i>Triticum aestivum</i> L. subsp. <i>aestivum</i> (*6)	Wheat	1,5	0,3	TR	80			5 000	250 000	500	25 000	30 000
<i>Triticum durum</i> – see <i>Triticum turgidum</i>												
<i>Triticum turgidum</i> L. subsp. <i>durum</i> (Desf.) van Slangeren [syn. <i>Triticum durum</i> Desf.] (*6)	Durum wheat	1	0,3	TR	80			5 000	250 000	500	25 000	30 000
<i>Vicia faba</i> L. (*3)	Broad bean	4	0,5	0,2	70			5 000	5 000	500	500	30 000
<i>Vicia sativa</i> L. [incl. <i>V. angustifolia</i> L.] (*3)	Common vetch	4	0,2		60			5 000	95 000 300 000	500	9 500 30 000	30 000
<i>Vicia villosa</i> Roth [incl. <i>V. dasycarpa</i> Ten.] (*3)	Hairy vetch	4	0,2		60			5 000	125 000 180 000	500	12 500 18 000	30 000
<i>Vigna unguiculata</i> (L.) Walp. [syn. <i>V. sinensis</i> (L.) Savi ex Hassk.; incl. <i>Dolichos biflorus</i> L.] (*3)	Cowpea	4	0,1		60			5 000	40 000	500	4 000	30 000
<i>Zea mays</i> L.	White and yellow maize	4	0,1		70			10 000	36 000	1 000	3 600	40 000
<i>Zea mays</i> L.	Sweetcorn	4	0,1		70			10 000	60 000	1 000	6 000	40 000

FOOTNOTE

1.	In the case of <i>Lespedeza cuneata</i> (Dum. Cours.) G. Don, <i>Kummerowia striata</i> (Thunb.) Schindl. [Syn.: <i>Lespedeza striata</i> (Thunb.) Hook. et Arn.], <i>Ornithopus compressus</i> L., <i>Stylosanthes hamata</i> (L.) Taub. and <i>Trifolium vesiculosum</i> Savi, up to 60 % hard seeds may be counted as germinated or viable seeds.
2.	In the case of <i>Desmodium intortum</i> (Mill.) Urb., <i>Desmodium uncinatum</i> (Jacq.) DC., <i>Medicago littoralis</i> Rohde ex Loisel., <i>Medicago polymorpha</i> L., <i>Medicago rugosa</i> Desr., <i>Medicago sativa</i> L., <i>Medicago scutellata</i> (L.) Miller, <i>Medicago truncatula</i> Gaertn. and <i>Trifolium repens</i> L., up to 40 % hard seeds may be counted as germinated or viable seeds.
3.	In the case of <i>Lotus corniculatus</i> L., <i>Lupinus albus</i> L., <i>Lupinus angustifolius</i> L., <i>Lupinus luteus</i> L., <i>Ornithopus sativus</i> Brot., <i>Trifolium fragiferum</i> L., <i>Trifolium hirtum</i> All., <i>Trifolium incarnatum</i> L., <i>Trifolium pratense</i> L., <i>Trifolium resupinatum</i> L., <i>Trifolium subterraneum</i> L., <i>Vicia faba</i> L., <i>Vicia sativa</i> L., <i>Vicia villosa</i> Roth and <i>Vigna unguiculata</i> (L.) Walp., [syn. <i>V. sinensis</i> (L.) Savi ex Hassk.; incl. <i>Dolichos biflorus</i> L.] up to 20 % hard seeds may be counted as germinated or viable seeds.
4.	In the case of <i>Sorghum</i> spp. the test for viability is only applicable where seed is still enveloped by the flower parts.
5.	In the case of <i>Eragrostis curvula</i> (Schrud.) Nees, not more than 0,2 % nematode galls caused by <i>Anguina</i> spp., are allowed.
6.	In the case of <i>Avena nuda</i> L., <i>Hordeum vulgare</i> L. subsp. <i>vulgare</i> , <i>Triticum aestivum</i> L. subsp. <i>aestivum</i> and <i>Triticum turgidum</i> L. subsp. <i>durum</i> (Desf.) van Slangeren [syn. <i>Triticum durum</i> Desf.], TR means trace components amounting to less than 0,05 %.
7.	In the case of <i>Anthephora pubescens</i> Nees and <i>Sporobolus fimbriatus</i> (Trin.) Nees, seed sampling and testing methods are available from the Official Seed Testing Laboratory.

[Table 4 amended, corrected and substituted by R. 1287 of 14 June 1985, R. 1524 of 12 July 1985, R. 1487 of 11 July 1986, R. 2496 of 9 December 1988, R. 1638 of 12 July 1991, R. 1971 of 16 August 1991, R. 2119 of 24 July 1992, R. 2618 of 18 September 1992, R. 1590 of 27 August 1993, R. 2057 of 29 October 1993, R. 513 of 18 March 1994, R. 174 of 10 February 1995, R. 319 of 3 March 1995, R. 97 of 24 January 1997, R. 1207 of 1 December 2000; R. 849 of 2 September 2005 (as corrected by R. 928 of 30 September 2005, R. 770 of 4 August 2006, R. 56 of 2 February 2007 and No. 1210 of 13 November 2020.)]

[Tabel 4 gewysig, verbeter en vervang deur R. 1287 van 14 Junie 1985, R. 1524 van 12 Julie 1985, R. 1487 van 11 Julie 1986, R. 2496 van 9 Desember 1988, R. 1638 van 12 Julie 1991, R. 1971 van 16 Augustus 1991, R. 2119 van 24 Julie 1992, R. 2618 van 18 September 1992, R. 1590 van 27 Augustus 1993, R. 2057 van 29 Oktober 1993, R. 513 van 18 Maart 1994, R. 174 van 10 Februarie 1995, R. 319 van 3 Maart 1995, R. 97 van 24 Januarie 1997, R. 1207 van 1 Desember 2000; R. 849 van 2 September 2005 (soos verbeter deur R. 928 van 30 September 2005), R. 770 van 4 Augustus 2006, R. 56 van 2 Februarie 2007 en No. 1210 van 13 November 2020.)]